



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

Multiple Integrals Over Infinite Fields, and the Fourier Multiple Integral.*

BY BURTON H. CAMP.

INTRODUCTION.

1. Multiple integrals over infinite fields have been defined in recent years in different ways by Pierpont,[†] Vallée-Poussin,[‡] and Hardy. § The integrals of Pierpont and of Vallée-Poussin exist only when they exist absolutely, but this is not the case with Hardy's integral. It is shown here that, for functions whose integrals over finite fields exist, these three infinite integrals are equivalent when any one of them exists absolutely. This paper is divided into three parts. In the first part (§§ 2-5) are developed relations between Hardy's integral and the iterated integrals, certain new theorems on inversion of the order of integration, and some fundamental properties of multiple infinite integrals, most of which are known for the case of integrals over finite fields but not for the infinite case. Among these last is the second theorem of the mean. Heretofore, this has not been proved, apparently, in the infinite case even for functions of one variable. The second part (§§ 6-8) deals with the behavior of integrals with respect to parameters. I am mainly concerned here with the infinite integrals, but it is necessary incidentally to extend to the case of integrals over finite fields in space of several dimensions theorems which are known only for integrals over one-dimensional fields. To save space, and because novel methods would not be introduced, the proofs of many of the statements in this part are not given. They have to do, however, with fundamental matters, including what are, in one dimension, well-known tests for termwise integration and differentiation; in the infinite case they include the so-called fundamental theorem of the integral calculus. A résumé of the third part (§§ 9-11), which contains certain applications to Fourier's double and quadruple integrals, is given at the beginning of § 9.

* Read in part before the American Mathematical Society, February 27, 1915, April 29, 1916.

[†] *Transactions of the American Mathematical Society*, Vol. VII (1906), p. 169.

[‡] *Cours d'Analyse*, 2d ed., Vol. II, pp. 69, 70, 108, 109.

§ *Messenger of Mathematics*, Vol. XXXII (1902-3), pp. 92 ff.

FUNDAMENTAL THEORY, §§ 2-5.

2. **Elementary Theorems.** As we shall not be concerned usually with integration over finite fields, we make the assumption always that the functions used are *absolutely L-integrable** in each *limited* measurable set of their domain of definition. Let R denote the infinite rectangle† where $x, y \geq 0$, and $R_{\alpha, \beta}$ the finite rectangle where $0 \leq x \leq \alpha$, $0 \leq y \leq \beta$. Let $f(x, y)$ be defined in R . We may then define the following integrals, when the limits in question exist:

$$J_1(f) = \lim_{\alpha, \beta = \infty} \int_{R_{\alpha, \beta}} f dx dy,$$

$$J_2(f) = \lim_{\alpha = \infty} \lim_{\beta = \infty} \int_{R_{\alpha, \beta}} f dx dy,$$

$$J_3(f) = \lim_{\beta = \infty} \lim_{\alpha = \infty} \int_{R_{\alpha, \beta}} f dx dy,$$

$$J_4(f) = \lim_{\alpha = \infty} \int_0^\alpha dx \lim_{\beta = \infty} \int_0^\beta f dy = \int_0^\infty dx \int_0^\infty f dy,$$

$$J_5(f) = \lim_{\beta = \infty} \int_0^\beta dy \lim_{\alpha = \infty} \int_0^\alpha f dx = \int_0^\infty dy \int_0^\infty f dx.$$

The integral $J_1(f)$ is Hardy's definition referred to in the introduction, J_4 and J_5 are the usual iterated integrals, J_2 and J_3 are introduced chiefly for convenience in developing the theory of J_1 , J_4 and J_5 . When it is desired to integrate a function over a larger portion or over all of the infinite two-dimensional field, we let R_1 denote the rectangle $(x, y \geq 0)$, R_2 the rectangle $(x < 0 \leq y)$, R_3 the rectangle $(x, y < 0)$, and R_4 the rectangle $(y < 0 \leq x)$, and let $R = R_1 + R_2 + R_3 + R_4$, and then define the integral over R as the sum of the integrals over R_1 , R_2 , R_3 , and R_4 . These definitions could be generalized slightly by using the concept of the Harnack-Lebesgue integral, but that is not worth while for my present purpose. By $\lim_{\alpha, \beta = \infty}$ is meant the double limit‡ as α, β diverge to plus infinity passing through all positive values. Most of the theorems, though stated only for two dimensions, may obviously be extended to n dimensions.

The two following theorems are almost immediate consequences of our definition, and the succeeding lemma is known.

* *I. e.*, integrable in the sense of Lebesgue.

† The case of n variables is not essentially different from the case of two, but the treatment would be cumbersome. In §6, however, where finite fields only are employed, and there is no loss of simplicity, n variables are used.

‡ *E. g.*, Pierpont, *Theory of Functions of Real Variables*, Vol. I (1905), p. 195.

THEOREM 1.* *A necessary and sufficient condition that $J_1(f)$ shall exist is that, for every pair of constants $h, k > 0$,*

$$\lim_{\alpha, \beta = \infty} \int_r f dx dy = 0,$$

where $r = R_{\alpha+h, \beta+k} - R_{\alpha, \beta}$.

THEOREM 2. *If $J_i(f)$ and $J_i(g)$ exist, and c is a constant, $J_i(f) + J_i(g) = J_i(f+g)$, and $J_i(cf) = cJ_i(f)$.*

LEMMA. *If for all values of $\Delta\alpha, \Delta\beta \geq 0$, $\phi(\alpha + \Delta\alpha, \beta + \Delta\beta) \geq \phi(\alpha, \beta)$, and $\phi(\alpha, \beta)$ is less than a fixed number M , then $\lim_{\alpha, \beta} \phi(\alpha, \beta)$, $\lim_{\alpha} \lim_{\beta} \phi(\alpha, \beta)$, $\lim_{\beta} \lim_{\alpha} \phi(\alpha, \beta)$ all exist and are equal and not greater than M .*

THEOREM 3. *If, in general, $\dagger 0 \leq f(x, y) \leq g(x, y)$, and if $J_i(g)$ exists, so does $J_i(f)$, and $J_i(f) \leq J_i(g)$.*

If $i=1, 2$, or 3 , this is an immediate consequence of the lemma. If $i=4$, by hypothesis there exists, except perhaps for a null set, a positive L -integrable function,

$$L(x) = \int_0^{\infty} g dy,$$

whose integral from 0 to α is a monotonic increasing function of α and has the limit $J_4(g)$. Except perhaps for a null set of x 's,

$$\int_0^{\beta} f dy \leq \int_0^{\beta} g dy,$$

and the former is a monotonic increasing function of β whose limit $l(x)$ is less than or at most equal to $L(x)$. The integral of $l(x)$ from 0 to α is a monotonic increasing function of α and hence

$$\lim_{\alpha} \int_0^{\alpha} l(x) dx = J_4(f) \leq \lim_{\alpha} \int_0^{\alpha} L(x) dx = J_4(g).$$

COROLLARY.‡ *The function $J_i(f)$ exists if $J_i(g)$ does, when $f(x, y) = g(x, y)$ in a part of R and zero elsewhere, provided that the field is so divided that f is L -integrable over every limited, measurable set in each part.*

THEOREM 4. *A necessary and sufficient condition that $J_i(f)$ shall exist is that there shall exist two other functions $g(x, y)$, $G(x, y)$, for both of*

*The proof of this depends on the following theorem, well known for functions of one variable: A necessary and sufficient condition that the double limit of $\phi(\alpha, \beta)$ shall exist, as α, β become infinite, is that, for every pair of constants $h, k > 0$, $\lim_{\alpha, \beta = \infty} [\phi(\alpha + h, \beta + k) - \phi(\alpha, \beta)] = 0$.

†I. e., except perhaps at a set of points of measure zero. Such a set will be called a null set.

‡This is not necessarily true if g has both signs. Indeed, this property, together with the properties in Theorem 2, would be sufficient to show that $|f|$ would be integrable, no matter what the definition of integration for f .

which J_i exists, such that in general $g(x, y) < f(x, y) < G(x, y)$. Then also $J_i(g) \leq J_i(f) \leq J_i(G)$.

Since $0 < f - g < G - g$, $J_i(f - g) \leq J_i(G - g)$. Then, since $J_i(g)$ exists, we have $J_i(f) \leq J_i(G)$.

COROLLARY 1. *The same is true, with an appropriate change in the last statement, if $g \leq f \leq G$.*

COROLLARY 2. *The function $J_i(f)$ exists if it exists absolutely, or if f is the product of two positive functions which are less than unity and for each of which J_i exists, or if it is the p -th power ($p > 1$) of such a function.*

COROLLARY 3. *Under the usual conditions the mean-value theorem, $mJ_i(g) \leq J_i(fg) \leq MJ_i(g)$, is valid.*

COROLLARY 4. *If $J_i(f^2)$ and $J_i(g^2)$ exist, then $J_i(fg)$ exists absolutely, and the inequality of Schwarz, $J_i^2(fg) \leq J_i(f^2) \cdot J_i(g^2)$, is valid.**

CASE I: ($i = 1$). Let $\psi = f$ and $\phi = 0$, $\gamma = g$ and $\delta = 0$ where $fg \geq 0$, and let $\phi = f$ and $\psi = 0$, $\delta = g$ and $\gamma = 0$ where $fg < 0$. The same equalities hold when both sides are squared, and by the corollary of Theorem 3 the following exist: $J_1(\psi^2)$, $J_1(\phi^2)$, $J_1(\gamma^2)$, $J_1(\delta^2)$. By the inequality for finite fields,

$$\left(\int_{R_{\alpha, \beta}} \psi \gamma dx dy \right)^2 \leq \left(\int_{R_{\alpha, \beta}} \psi^2 dx dy \right) \left(\int_{R_{\alpha, \beta}} \gamma^2 dx dy \right) \leq J_1(\psi^2) \cdot J_1(\gamma^2).$$

Since $\psi \gamma \geq 0$, the function of α, β on the left satisfies the conditions of the lemma to Theorem 3. Therefore $J_1^2(\psi \gamma)$ exists, and so does $J_1(\psi \gamma)$. Similarly, $J_1(\phi \delta)$ exists, and since by the early equalities $|fg| = \psi \gamma - \phi \delta$, so does $J_1(|fg|)$. Now, proceeding to the limit in the relation,

$$\left(\int_{R_{\alpha, \beta}} fg dx dy \right)^2 \leq \left(\int_{R_{\alpha, \beta}} f^2 dx dy \right) \left(\int_{R_{\alpha, \beta}} g^2 dx dy \right),$$

we may establish the proposition.

CASE II: ($i > 1$). This may be referred to Case I by the use of the corollary to Theorem 5 and the first part of Corollary 2, Theorem 4.

3. **The Integrals of Pierpont and Vallée-Poussin.**[†] In considering these definitions we shall assume as in § 2 that $f(x, y)$ is absolutely L -integrable in every limited measurable part of the field R . This is not done by Pierpont, but it is better for our present purpose to dispense with the extra element of generality which he obtains by omitting this condition. Let R be defined as before, and let $D_1 < D_2 < \dots$ be any infinite sequence of limited, measurable

*It does not follow, as in the finite case, that f is integrable if f^2 is.

† Cf. § 1.

point sets of such a nature that each D is wholly in R and that every point of R is in some D .

Vallée-Poussin's Definition, $V(f)$. Let $f \geq 0$. If the following limit exists and is independent of the choice of D 's,

$$V(f) = \int_R f dx dy = \lim_{i=\infty} \int_{D_i} f dx dy. \quad (1)$$

Let f be unrestricted as to sign. Set $f = f_1 - f_2$, where $f_1, f_2 \geq 0$. When the integrals on the right exist,

$$V(f) = \int_R f dx dy = \int_R f_1 dx dy - \int_R f_2 dx dy. \quad (2)$$

Pierpont's Definition, $P(f)$. Let f be unrestricted as to sign. Let the following be an infinite sequence of limited measurable sets, E_1, E_2, \dots ; and let E_i contain every point whose distance from the origin is less than i ; it may contain other points. When the following limit exists and is independent of the choice of E 's,

$$P(f) = \int_R f dx dy = \lim_{i=\infty} \int_{E_i} f dx dy. \quad (3)$$

It will be shown that this definition is equivalent to the definition:

$$P(f) = \lim_{i=\infty} \int_{D_i} f dx dy. \quad (4)$$

It is obvious that if (3) exists (4) exists and gives the same result. To prove the converse we let (4) exist. Suppose that, for a certain E set, the limit in (3) should not exist, *i. e.*, suppose that the sequence

$$\int_{E_1} f dx dy, \int_{E_2} f dx dy, \dots \quad (5)$$

should diverge. In this sequence there would exist a divergent sub-sequence

$$\int_{E'} f dx dy, \int_{E''} f dx dy, \dots \quad (6)$$

for which $E' < E'' < \dots$. For example, we might take $E' = E_1$, and then, since E_1 is limited, there would exist an E_i which would include all points as near to the origin as any of those of E_1 ; let $E'' = E_i$, or some E farther on in the sequence, if necessary, in order to make (6) diverge. This sub-sequence would be a D sequence.

LEMMA. If $f(x, y) \geq 0$ and the following limit exists for one set of D 's, it does for every set:

$$\lim_{i=\infty} \int_{D_i} f dx dy.$$

THEOREM 5. (a) If $f \geq 0$, $V(f) = P(f) = J_1(f)$, provided one of these exists. (b) If $V(f)$ or $P(f)$ exists, it exists absolutely, and then $V(f) = P(f) = J_1(f)$. (c) $J_1(f)$ may exist when $V(f)$ and $P(f)$ do not.

(a) Since $R_{\alpha, \beta}$ is a special case of E_i , if P exists, so does J_1 , and $J_1 = P$. By the lemma, if J_1 exists, so does V , and $V = J_1$. By definition $V = P$.

(b) It is known and obvious that if $V(f)$ exists so does $V(|f|)$, and

$$V(|f|) = \lim_{i=\infty} \int_{D_i} f_1 dx dy + \lim_{i=\infty} \int_{D_i} f_2 dx dy, \quad V(f) = \lim_{i=\infty} \int_{D_i} f_1 dx dy - \lim_{i=\infty} \int_{D_i} f_2 dx dy,$$

where $|f| = f_1 + f_2$, $f = f_1 - f_2$. By (a), then, $J_1(f_1) = V(f_1)$, $J_1(f_2) = V(f_2)$. Therefore, $J_1(f) = J_1(f_1) - J_1(f_2) = V(f)$. A similar result may be obtained for $P(f)$, since it may be shown that this integral also obeys Theorem 2. Thus (b) is proved if the hypothesis be that $V(f)$ exists. It only remains to show that if $P(f)$ exists so does $V(f)$. The following equation is true if two of the limits exist:

$$P(f) = \lim_{i=\infty} \int_{D_i} f dx dy = \lim_{i=\infty} \int_{D_i} f_1 dx dy - \lim_{i=\infty} \int_{D_i} f_2 dx dy.$$

If it is true, $V(f)$ exists. If it is not true, but $P(f)$ exists; since $f_1, f_2 \geq 0$, both of the last two integrals must diverge to plus infinity for some set of D 's, and therefore by the lemma for all sets. On this hypothesis we will show that there exists a set of D 's for which the first limit does not exist. Let R_1 be a circle of unit radius about the origin. Let

$$k_1 = \int_{R_1} f dx dy.$$

Since the integral of f_1 over R_1 is not infinite, and since by definition $f_1 = f$ where $f \geq 0$, there must exist outside of R_1 a limited set A_1 , in which $f \geq 0$, such that

$$\int_{A_1} f_1 dx dy > 1 - k_1.$$

Hence,

$$\int_{R_1 + A_1} f dx dy = \int_{R_1} f dx dy + \int_{A_1} f dx dy > k_1 + 1 - k_1 = 1.$$

Let $D_1 = R_1 + A_1$. Let R_2 be a circle about the origin including all of D_1 . Let

$$k_2 = \int_{R_2} f dx dy.$$

The integral of f_1 over R_2 is not infinite, and therefore, as before, there exists a set A_2 outside of R_2 where $f \geq 0$ such that

$$\int_{R_2 + A_2} f dx dy = \int_{R_2} f dx dy + \int_{A_2} f dx dy = \int_{R_2} f dx dy + \int_{A_2} f_1 dx dy > 2.$$

Let $D_2 = R_2 + A_2$; etc.

(c) Hardy* has given some interesting examples of this. One which will concern us later is $J_1(\cos xy) = \pi/2$.

COROLLARY.† If $J_i(|f|)$ exists, so does $J_j(|f|)$, and $J_i(|f|) = J_j(|f|)$, and $J_i(f) = J_j(f)$.

Vallée-Poussin has shown that if $V(|f|)$ exists, $J_4(|f|) = J_5(|f|) = V(|f|)$, and that $J_4(f) = J_5(f) = V(f)$. By the theorem, then, the corollary is proved for the cases $i, j = 1, 4, 5$. The remaining cases may easily be proved by the use of the lemma to Theorem 3.

4. **More Existence and Inversion Theorems.** THEOREM 6. Let $J_i(f)$ exist. A necessary and sufficient condition that $J_j(f)$ shall exist is that there shall exist a function $g(x, y)$ which is in general less than $f(x, y)$ for which both J 's exist.

Necessity. Let $q(x, y) > 0$, and let $J_j(q)$ exist. We may take $g = f - q$, for, since $J_i(f)$ and $J_j(q)$ exist, so does $J_j(f - q)$.

Sufficiency. Since $J_i(f - g)$ exists and $f - g \geq 0$, $J_j(f - g)$ exists by the preceding corollary. Since now $J_j(g)$ exists, so does $J_j(f - g + g)$.

COROLLARY 1. Let $J_i(f)$ exist. A necessary and sufficient condition that $J_j(f) = J_i(f)$ is that there shall exist a function $g(x, y)$, in general less than $f(x, y)$, for which this is true.

$J_i(f - g) = J_j(f - g) = J_i(f) - J_i(g)$. Add the equality, $J_i(g) = J_j(g)$.

COROLLARY 2. Let one of the iterated integrals $J_4(f)$, $J_5(f)$ exist. A necessary and sufficient condition that inversion of the order of iteration be allowable is that there shall exist a function $g(x, y)$, in general less than $f(x, y)$, for which the inversion is allowable.

COROLLARY 3.‡ A necessary and sufficient condition that both iterated integrals of f shall exist and be equal is that this shall be true of two functions $g(x, y)$, $G(x, y)$ for which in general $g < f < G$.

*Loc. cit., § 1, pp. 159 ff.

† Vallée-Poussin, loc. cit., Vol. II, p. 121. Bromwich (*Proceedings of the London Mathematical Society*, Vol. I (1903), p. 181, third footnote) indicates that this result follows from the lemma to Theorem 3, but really only the last part, in which it is said that $J_1 = J_2 = J_3$, follows from that lemma. Indeed, if it did follow also that $J_1 = J_4 = J_5$, as he states, his result would antedate its discovery for the finite case by four years (see also Hobson, same *Proceedings*, Vol. VIII (1910), p. 22).

I am omitting from this paper the subject of transformation of variables in these integrals. It is true, however, that, if any J_i exists absolutely, it may be transformed into an equivalent double, or iterated, integral over the unit square. Another proof of the corollary may be devised by the use of transformations of this sort.

‡ Cf. W. H. Young, same *Proceedings*, Vol. IX (1910-11), p. 323. Young's theorem gives only sufficient criteria that inversion of the order of iteration be permissible, and involves the additional hypothesis that this inversion be permissible for integrals of g and G over one-way infinite fields.

COROLLARY 4.* (a) *Inversion of the order of iteration of f over the one-way infinite field $R_\alpha(0 \leq x \leq \alpha, 0 \leq y)$ is allowable if one of the iterated integrals of $|f|$ exists.* (b) *If one of the iterated integrals of f over R_α exists, a necessary and sufficient condition that inversion be allowable is that there shall exist a function $g(x, y)$, in general less than $f(x, y)$, for which it is allowable.* (c) *A necessary and sufficient condition that both iterated integrals over R_α shall exist and be equal is that there shall exist two functions $g(x, y)$, $G(x, y)$ for which this is so and for which in general $g < f < G$.*

COROLLARY 5. *The theorem and previous corollaries are true if we write, instead of $g < f (f < G)$, $g \leq f (f \leq G)$, or if, in the theorem and first two corollaries, we substitute $G > f$ for $g < f$.*

THEOREM 7. (a) *If $J_1(f)$ exists, and $J_2(f)$ exists over R_α for sufficiently large values of α (say $\alpha > C$) then $J_2(f) = J_1(f)$; similar statements may be made for J_3 and the one-way infinite field $R_\beta(0 \leq y \leq \beta, 0 \leq x)$.* (b) *Let inversion[†] of the order of iteration over $R_\alpha(\alpha > C)$ be allowable; then $J_2(f) = J_4(f)$ if either exists; and similar statements may be made for J_3 and J_5 and R_β .*

COROLLARY. *Sufficient conditions that all the J 's shall exist and be equal are that inversion be allowable over both R_α and $R_\beta(\alpha, \beta > C)$, and that J_1 and one of each of the pairs (J_2, J_4) , (J_3, J_5) shall exist. Then also all the J 's are equal.*

The proofs of the theorems and corollaries of this section are simple.

5. **Second Theorem of the Mean.**[‡] THEOREM 8. *Let $f(x, y)$ be limited in R and L -integrable over every finite measurable set, and let $g(x, y)$ be absolutely L -integrable in R . Then, if $l \leq f(x, y) \leq L$, there exists a k such that*

$$J = \int_R f(x, y) g(x, y) dx dy = l \int_{R-A_k} g dx dy + L \int_{A_k} g dx dy;$$

where, if f has the property that the set of points E_k , where $f = k$, is always null, A_k is defined as the set where $f > k$; and otherwise A_k is the set where

* Parts (a) and (c) are known. See also note to Theorem 7 (b). By the existence of the integral of f over R_α we mean the existence over R of the integral of the function which equals f in R_α and zero elsewhere.

† For criteria for this condition we have, not only Theorem 6, Corollary 4, but also the usual condition that, assuming $|f|$ L -integrable over finite fields, termwise integration with respect to x of the β sequence—viz: the integral of f with respect to y from 0 to β —be allowable in $(0, \alpha)$. This assumes implicitly the existence of one iterated integral.

‡ Proved for the case where R is finite by Lebesgue, *Annales de l'École Normale*, (3) Vol. XXVII (1910), p. 444; proved in a slightly more general form by the author, *Mathematische Annalen*, Vol. LXXV (1914), p. 285. There will be used in this section much of the notation and some of the results of the latter paper.

$f > k$, plus perhaps certain points where $f = k$. In the second case the set A_k may be defined by the use of an associated function $q(x, y)$.

CASE 1: ($l=0, \hat{E}_k=0$).^{*} Consider a monotonic sequence of circles whose union[†] is $R: R_1 < R_2 < \dots$. Let A_k be the set of points where $f > k$. Let A'_{k_i} refer to the parts of A_{k_i} in R_i . By the finite case there exists for each R_i a k_i such that

$$\int_{R_i} fg = L \int_{A'_{k_i}} g. \quad (1)$$

By Theorem 4, fg is absolutely L -integrable in R , and by Theorem 5

$$J = \int_R fg = \lim_{i \rightarrow \infty} \int_{R_i} fg = L \lim_{i \rightarrow \infty} \int_{A'_{k_i}} g; \quad (2)$$

so the last limit exists. It only remains to show that there is a definite k and A_k for which

$$\lim_{i \rightarrow \infty} \int_{A'_{k_i}} g = \int_{A_k} g. \quad (3)$$

The left-hand side may be considered as the limit of any infinite sub-sequence selected from the sequence

$$\int_{A'_{k_1}} g, \int_{A'_{k_2}} g, \dots \quad (4)$$

Suppose the sub-sequence to be the terms corresponding to $i = a_1 < a_2 < \dots$, where these numbers are so chosen that, if k be taken as the lower limit of the sequence k_1, k_2, \dots , it is also the limit of either (5) or (6),

$$k_{a_1} \geq k_{a_2} \geq \dots, \quad (5)$$

$$k_{a_1} \leq k_{a_2} \leq \dots \quad (6)$$

To be assured that this is possible we first note that whenever the points k_1, k_2, \dots constitute an infinite point set, there exists a lower limit, for the set is limited ($0 \leq k_i \leq L$). Whenever the points do not constitute an infinite point set, certain of these numbers are repeated an infinite number of times; let k be the least of these numbers. It is immaterial to the proof whether (5) or (6) holds. Assume (5); and to avoid the additional subscripts let it be assumed that (4) is the sequence corresponding to (5), *i. e.*, that k is the limit of k_i , and that

$$k_1 \geq k_2 \geq \dots \quad (7)$$

Then

$$A'_{k_1} \leq A'_{k_2} \leq \dots \quad (8)$$

^{*} By \hat{E} is meant the measure of E , a concept which may be defined in the infinite case as the integral over E of unity.

[†] By "union" of (R_i) is meant the totality of points each of which belongs to some R_i .

The limit of this sequence may be an infinite field. For $\varepsilon > 0$ prescribed, let R_a be so large that

$$\int_{R-R_a} |g| < \varepsilon. \quad (9)$$

Let

$$A''_{k_1} \subseteq A''_{k_2} \subseteq \dots \quad (10)$$

refer to the portions of A'_{k_i} , etc., in R_a . If $i < \alpha$, A''_{k_i} may not refer to all of the points of R_a where $f > k_i$, for A'_{k_i} may not include them all; but if $i \geq \alpha$, A''_{k_i} does include all those points, and only those. A''_k is the set in R_a where $f > k$; and so, by the reasoning of my *Annalen* paper,* A''_k is the union of A''_{k_i} and $\lim_{i=\infty} \hat{A}''_{k_i} = \hat{A}''_k$. Thus, on account of the absolute continuity of the integral of g in R_a , there exists an $i_\varepsilon > \alpha$ and so large that if $i > i_\varepsilon$ ($\hat{A}''_k - \hat{A}''_{k_i}$) is sufficiently small to make

$$\int_{A''_k - A''_{k_i}} |g| < \varepsilon. \quad (11)$$

Now the set $(A_k - A'_{k_i})$ is made up of two parts, $(A''_k - A''_{k_i})$ and B , where B is some set outside of R_a . Therefore, by (9) and (11),

$$\left| \int_{A_k - A'_{k_i}} g \right| < \int_{A''_k - A''_{k_i}} |g| + \int_B |g| < 2\varepsilon,$$

which proves (3), and hence the proposition.

CASE 2: ($l=0$, $\hat{E}_k \geq 0$). The only new difficulty here is with inequality (11). It is obvious that $(\hat{A}''_k - \hat{A}''_{k_i})$ may not become small at pleasure because of the non-null sets that may be included. Therefore a method must be found of dividing up these non-null E_k sets. The measures of some of them may be infinite. We first note that there are only an enumerable number of values of k , say k_1, k_2, \dots , where $\hat{E}_k > 0$. This follows from the fact that, if A_k be, as before, the set where $f > k$, \hat{A}_k is a monotonic decreasing function of k , and k_i is a point of discontinuity of \hat{A}_k . It is known that such points are at most enumerably infinite for monotonic functions. We now define, corresponding to the parameter λ ($0 \leq \lambda \leq L+1$), a family of point sets Ω_λ , and a limited function $q(x, y)$ which is associated in a certain way with $f(x, y)$, and which is constant at most in null sets, and has the property that the set of points where $q > \lambda$ is Ω_λ . Let c_i be any positive number such that

$$c_1 + c_2 + \dots = 1. \quad (12)$$

**Loc. cit.*, p. 275, Lemma 2.

Let $q(x, y)$ be defined as the limit of the monotonic sequence,

$$f(x, y) \leq q_1(x, y) \leq q_2(x, y) \leq \dots \leq f(x, y) + 1, \quad (13)$$

where q_1, q_2, \dots are defined as follows: at points where $f < k_1$, let q_1 equal f ; at points where $f = k_1$, let q_1 equal $c_1 e^{-x^2-y^2} + f$; and at points where $f > k_1$, let q_1 equal $f + c_1$. Here the function $c_1 e^{-x^2-y^2} + k_1$ has the properties that it is constant only in null sets, that it lies between k_1 and $k_1 + c_1$, and that the set of points where it is greater than any number between these limits is a circle. Other definitions of q_1 might obviously be chosen, resulting in other associated functions q . Next, let q_2 be equal to q_1 where $f < k_2$, equal to $c_2 e^{-x^2-y^2} + q_1$ where $f = k_2$, and equal to $q_1 + c_2$ where $f > k_2$, etc. Let Ω_λ be the set of points where $q(x, y) > \lambda$ ($0 \leq \lambda \leq L+1$). The proof that q is constant at most in null sets is omitted because the fact seems obvious.

It will now be shown that each Ω_λ is divisible into two parts; one is part or all of the set where f equals a certain corresponding k_λ , and the other is the whole of the set where $f > k_\lambda$. Consider the set A_k where $f > k$, and suppose the k_i 's (7) which are not greater than k to be denoted by α_i 's and their corresponding c_i 's by s_i 's, thus:

$$\alpha_1, \alpha_2, \dots \leq k; \quad s_1, s_2, \dots$$

In A_k ,

$$q \geq f + s_1 + s_2 + \dots > k + s_1 + s_2 + \dots \quad (14)$$

For example, if $k_1 = \alpha_1, k_2 > k, k_3 = \alpha_2, \dots$; in $A_k, q_1 = f + s_1, q_2 \geq q_1 = f + s_1, q_3 = q_2 + s_2 \geq f + s_1 + s_2$, etc., and finally $q_1 \leq q_2 \leq \dots \leq q$. In $R - A_k$,

$$q \leq f + s_1 + s_2 + \dots \leq k + s_1 + s_2 + \dots, \quad (15)$$

for here, for every $j, q_j \leq f + s_1 + s_2 + \dots$, and finally $q = \lim_j q_j$. Thus, if $\lambda = k + s_1 + s_2 + \dots, q > \lambda$ in A_k and $q \leq \lambda$ in $R - A_k$, and so in this case $\Omega_\lambda = A_k$. If now it could be shown that to every λ there corresponds a k such that $\lambda = k + s_1 + s_2 + \dots$, the part of our proposition considered in this paragraph would be established, but this is not always true. It can, however, be proved except for an enumerable number of intervals where

$$\lambda_{k_i} - c_i < \lambda \leq \lambda_{k_i}. \quad (16)$$

For $s_1 + s_2 + \dots$ is a monotonic increasing function of k , and therefore, considered as a function of $k, \lambda = k + s_1 + s_2 + \dots$ is always increasing, and is defined for each k in $(0, L)$. Moreover, it is always continuous on the right, for, for any \bar{k} which is not a k_i , there exists a δ so that the interval $(k, k + \delta)$ does not contain any of the points k_1, k_2, \dots, k_n (n finite). Therefore,

$$|\lambda(\bar{k} + \delta) - \lambda(\bar{k})| < |\delta| + \sum_n^{\infty} c_i < \varepsilon,$$

if n and δ are chosen properly. This is true for $\delta > 0$ even if \bar{k} is a k_i , but in this case every interval $(\bar{k}, \bar{k} + \delta)$, $\delta < 0$, would include k_i , and hence at such points λ would have a discontinuity on the left equal to c_i . The correspondence desired is proved. We proceed to consider values of λ of (16). First, all points of A_{k_i} are in Ω_λ , for, if $f > k_i$, $q > \lambda_{k_i} = k_i + s_1 + s_2 + \dots$ ($c_i = s_1$), that is $q > \lambda$. Secondly, Ω_λ contains no other points, except some where $f = k_i$; for, if P is a point where $f(P) < k_i$, let $f(P) = k_i - \gamma$. To this value of k there is a corresponding $\bar{\lambda} = k_i - \gamma + \sigma_1 + \sigma_2 + \dots$, where $\sigma_1, \sigma_2, \dots$ are the s_i 's for $k_i - \gamma$; their sum is less than $s_1 + s_2 + \dots$ by at least $c_i = s_1$. By (15)

$$q(P) \leq k_i - \gamma + \sigma_1 + \sigma_2 + \dots \leq k_i - \gamma + s_1 + s_2 + \dots - s_1 < \lambda_{k_i} - c_i < \lambda,$$

and therefore P is not in Ω_λ . Thus, in every case there is a k_λ corresponding to λ for which the conditions are satisfied.

It may now be shown that

$$J = L \int_{\Omega_\lambda} g.$$

As in Case 1 (1),

$$\int_{R_i} fg = L \int_{\Omega'_{\lambda_i}} g,$$

where Ω'_{λ_i} is the part of Ω_{λ_i} in R_i . For q is for R_i precisely an associated function in the sense of my paper for the case when R is limited. The rest of the reasoning of Case 1 now holds, *mutatis mutandis*.

CASE 3: ($l \neq 0$). We may obviously consider Case 1 as a special case of Case 2, where $q = f$. By Case 2,

$$\int_R [f(x, y) - l]g(x, y) = (L - l) \int_{\Omega_\lambda} g = L \int_{\Omega_\lambda} g - l \int_R g + l \int_{R - \Omega_\lambda} g.$$

Add

$$\int_R lg = l \int_R g.$$

COROLLARY. If $f(x)$ is monotonic decreasing in (a, ∞) and $l \leq f(x) \leq L$, and if $g(x)$ is absolutely L -integrable in the same interval, there exists a λ ($a \leq \lambda \leq \infty$) such that

$$\int_a^\infty fg = l \int_\lambda^\infty g + L \int_a^\lambda g.$$

It is to be noted also that the theorem gives even for one dimension a much more general statement than this corollary.

INTEGRALS CONTAINING PARAMETERS, §§ 6–8.

6. **Integrals over Finite Fields.*** As a part of my discussion of integrals containing parameters I intend to extend to integrals over infinite fields certain theorems of W. H. Young† on integrals of functions of one variable. It is necessary first to make the extension to n dimensions for finite fields, and in this section I shall cite certain of these results. They may be deduced more simply from Vitali's test‡ of equi-absolute continuity, than by extending to n dimensions Young's idea of upper (lower) semi-continuity, but I will not give the proofs. In passing, however, I call attention to one theorem (11) which is not related to Young's, but which is a generalization of a very useful theorem on integrals not containing parameters.

THEOREM 9. *Let A be a limited measurable point set in space of n dimensions. Let $\lim_{n,m} f_{n,m}(x, y, \dots) = f(x, y, \dots)$ in A , in general. The necessary and sufficient condition that*

$$\lim_{n,m \rightarrow \infty} \int_A f_{n,m} dA = \int_A f dA$$

is that there shall exist two functions $g_{n,m}(x, y, \dots)$, $G_{n,m}(x, y, \dots)$ whose limits are $g(x, y, \dots)$ and $G(x, y, \dots)$, respectively, for which this is true, and such that, in general, $g_{n,m} < f_{n,m} < G_{n,m}$.

COROLLARY 1. *An analogous theorem is true for iterated integrals.*

COROLLARY 2. *If $f_{n,m}(x, y, \dots) \geq 0$, and its limit is $f(x, y, \dots)$, and if also the limit of its integral over A is the integral of f over A , then a similar statement is true for the limit of its integral over any measurable part of A .*

THEOREM 10. *If the limit of a monotonic sequence of integrable functions $0 \leq f_1(x, y, \dots) \leq f_2(x, y, \dots) \leq \dots$, is integrable in A , the sequence is integrable termwise.*

THEOREM 11.§ *If in A $f(x, y, \dots)$ is the limit of a function $f_{n,m}(x, y, \dots)$ which is nowhere negative, and the limit of whose integral is zero, then in general $f=0$.*

The proof is immediate if it can be shown that the integral of f over A exists, for in that case the integral of $f_{n,m}$ is obviously equi-absolutely con-

*It is not assumed in this section, as it is elsewhere, that the functions in question have integrals over finite measurable fields. The case of n parameters is not essentially different from that of two.

† *Loc. cit.*, pp. 315 ff.

‡ *Rendiconti del Circolo di Palermo*, Vol. XXIII (1907) pp. 137 ff. Vitali enunciated his test only for one dimension, but his method is valid for n dimensions.

§ This is a generalization of a well-known theorem. Cf. Pierpont, *The Theory of Functions of Real Variables*, Vol. II (1911), p. 385, § 402 (2).

tinuous and approaches the integral of f as a limit. Thus the integral of f is zero, and the theorem of which this is a generalization applies. Suppose the integral of f over A did not exist. We could then let A_k be the set where $f > k$, and set ϕ_k equal to zero in A_k and to f in $(A - A_k)$. Then the integral of ϕ_k over A would diverge to plus infinity as k became infinite. This would be impossible, for suppose that for each k $\phi_{n,m;k}$ equals 0 in A_k and $f_{n,m}$ elsewhere. Then for each k $\phi_{n,m;k}$ would satisfy the conditions of the theorem and ϕ_k would be integrable. Therefore the integral of ϕ_k over A would equal zero for each k .

7. **Continuity.** LEMMA 1. *If $0 \leq f(x, y) \leq g(x, y)$, and $J_i(g)$ exists, then*

$$0 \leq J_i(f) - \int_{R_{\alpha, \beta}} f dx dy \leq J_i(g) - \int_{R_{\alpha, \beta}} g dx dy.$$

LEMMA 2. *If for each (n, m) the function $f_{n,m}(x, y)$ satisfies the conditions imposed on f in Lemma 1, and if $f(x, y) = \lim_{n,m} f_{n,m}(x, y)$ in general, then $J_i(f_{n,m})$ exists and its limit is $J_i(f)$.*

Obviously $0 \leq f \leq g$, and so by Theorem 3 $J_i(f_{n,m})$ and $J_i(f)$ exist. For an arbitrary $\varepsilon > 0$, let α, β be chosen so large that

$$J_i(g) - \int_{R_{\alpha, \beta}} g < \varepsilon.$$

By Lemma 1, then,

$$0 \leq J_i(f) - \int_{R_{\alpha, \beta}} f \leq J_i(g) - \int_{R_{\alpha, \beta}} g < \varepsilon, \quad (1)$$

$$0 \leq J_i(f_{n,m}) - \int_{R_{\alpha, \beta}} f_{n,m} \leq J_i(g) - \int_{R_{\alpha, \beta}} g < \varepsilon. \quad (2)$$

By Theorem 9 there exists for these fixed values of α, β an n_ε so large that, if $n, m > n_\varepsilon$,

$$\left| \int_{R_{\alpha, \beta}} f - \int_{R_{\alpha, \beta}} f_{n,m} \right| < \varepsilon. \quad (3)$$

From (1), (2), and (3),

$$|J_i(f) - J_i(f_{n,m})| < 3\varepsilon, \quad (n, m > n_\varepsilon).$$

LEMMA 3. *Except perhaps for a null set in R , let $g_{n,m}(x, y) \geq 0$, $\lim_{n,m} g_{n,m}(x, y) = g(x, y)$, and $\lim_{n,m} J_i(g_{n,m}) = J_i(g)$. Then, for all values of α, β uniformly,*

$$\lim_{n,m} \int_{R_{\alpha, \beta}} g_{n,m} dx dy = \int_{R_{\alpha, \beta}} g dx dy.$$

Let $\phi_{n,m} = g_{n,m}$ and $\psi_{n,m} = g$ where $g_{n,m} \geq g$, and let $\phi_{n,m} = g$ and $\psi_{n,m} = g_{n,m}$ where $g_{n,m} < g$. Then $0 \leq \psi_{n,m} < \phi_{n,m}$,

$$0 \leq \phi_{n,m} - \psi_{n,m} = |g_{n,m} - g|, \quad (1)$$

$\lim_{n,m} \phi_{n,m} = \lim_{n,m} \psi_{n,m} = g$, and $0 \leq \psi_{n,m} \leq g$. By Lemma 2, then,

$$\lim_{n,m} J_i(\psi_{n,m}) = J_i(g). \quad (2)$$

By definition, $\phi_{n,m} + \psi_{n,m} = g_{n,m} + g$, and hence $\lim_{n,m} J_i(\phi_{n,m} + \psi_{n,m}) = 2J_i(g)$. This, with (1) and (2), proves that $\lim_{n,m} J_i(|g_{n,m} - g|) = 0$. Now, uniformly in α, β ,

$$\left| \int_{R_{\alpha,\beta}} (g_{n,m} - g) \right| \leq \int_{R_{\alpha,\beta}} |g_{n,m} - g| \leq J_1(|g_{n,m} - g|).$$

LEMMA 4. *Except perhaps for a null set in R , let $0 \leq f_{n,m}(x, y) \leq g_{n,m}(x, y)$, let $\lim_{n,m} f_{n,m}(x, y) = f(x, y)$, and $\lim_{n,m} g_{n,m}(x, y) = g(x, y)$, and $\lim_{n,m} J_i(g_{n,m}) = J_i(g)$. Then $J_i(f_{n,m})$ exists and its limit is $J_i(f)$.*

Since obviously $0 \leq f \leq g$, we have, by Lemma 1, for all values of $n, m; \alpha, \beta$,

$$0 \leq J_i(f) - \int_{R_{\alpha,\beta}} f \leq J_i(g) - \int_{R_{\alpha,\beta}} g, \quad (1)$$

$$0 \leq J_i(f_{n,m}) - \int_{R_{\alpha,\beta}} f_{n,m} \leq J_i(g_{n,m}) - \int_{R_{\alpha,\beta}} g_{n,m}. \quad (2)$$

For each $\epsilon > 0$ there exists a number n_ϵ so that, by hypothesis and by Lemma 3, for all values of $n, m > n_\epsilon$ and all values of α, β uniformly,

$$|J_i(g_{n,m}) - J_i(g)|, \quad \left| \int_{R_{\alpha,\beta}} g - \int_{R_{\alpha,\beta}} g_{n,m} \right| < \epsilon, \quad (3)$$

and so

$$\left| \left(J_i(g_{n,m}) - \int_{R_{\alpha,\beta}} g_{n,m} \right) - \left(J_i(g) - \int_{R_{\alpha,\beta}} g \right) \right| < 2\epsilon. \quad (4)$$

But, if $\alpha, \beta > \text{some number } \alpha_\epsilon$,

$$\left| J_i(g) - \int_{R_{\alpha,\beta}} g \right| < \epsilon, \quad (5)$$

and, therefore, by (3) and (5),

$$\left| J_i(g_{n,m}) - \int_{R_{\alpha,\beta}} g_{n,m} \right| < 3\epsilon, \quad (n, m > n_\epsilon; \alpha, \beta > \alpha_\epsilon).$$

This, with (1) and (2), shows that

$$\left| J_i(f) - J_i(f_{n,m}) + \int_{R_{\alpha,\beta}} f_{n,m} - \int_{R_{\alpha,\beta}} f \right| < 4\epsilon (n, m > n_\epsilon; \alpha, \beta > \alpha_\epsilon). \quad (6)$$

Now let α, β be held fast at some values greater than α_ϵ . There exists a number n'_ϵ so that

$$\left| \int_{R_{\alpha, \beta}} f_{n, m} - \int_{R_{\alpha, \beta}} f \right| < \epsilon, \quad (n, m > n'_\epsilon), \quad (7)$$

for, by Lemma 3, and since $0 \leq f_{n, m} \leq g_{n, m}$, Theorem 9 applies. Thus (6) and (7) prove the lemma.

THEOREM 12. *Let $\lim_{n, m} f_{n, m}(x, y) = f(x, y)$ in general. In order that the following integrals shall exist and that*

$$\lim_{n, m} J_i(f_{n, m}) = J_i(f),$$

it is necessary and sufficient that there shall exist two convergent sequences of functions, $\lim_{n, m} g_{n, m}(x, y) = g(x, y)$, $\lim_{n, m} G_{n, m}(x, y) = G(x, y)$, for which this is true, such that in general $g_{n, m}(x, y) < f_{n, m}(x, y) < G_{n, m}(x, y)$.

To prove the necessary part we may take $g_{n, m} = f_{n, m} - 1/(mnx^2y^2)$, $G_{n, m} = f_{n, m} + 1/(mnx^2y^2)$. To prove the sufficient part we note that $0 \leq f_{n, m} - g_{n, m} \leq G_{n, m} - g_{n, m}$, and that $\lim_{n, m} J_i(G_{n, m} - g_{n, m}) = J_i(G - g)$, and apply Lemma 4.

COROLLARY 1. *Let $\lim_{n, m} f_{n, m}(x, y) = f(x, y)$ in general. Any one of the following is sufficient to make $\lim_{n, m} J_i(f_{n, m}) = J_i(f)$:*

- (a) $g(x, y) \leq f_{n, m}(x, y) \leq G(x, y)$, and $J_i(g)$, $J_i(G)$ shall exist;
- (b) $f(x, y)$ shall be the limit of a monotonic sequence,

$$f_1(x, y) \leq f_2(x, y) \leq \dots,$$

and $J_i(f)$ and $J_i(f_1)$ shall exist;

- (c) $\lim_{n, m} J_i(|f_{n, m}|) = J_i(|f|)$;
- (d) $|f_{n, m}(x, y)| \leq G(x, y)$ and $J_i(G)$ shall exist;
- (e) $f_{n, m}(x, y) = h_{n, m}(x, y)g_{n, m}(x, y)$, where $0 \leq h_{n, m}, g_{n, m} \leq 1$, and one of the convergent sequences $h_{n, m}$ and $g_{n, m}$ shall be integrable termwise in R ;
- (f) $f_{n, m} = (g_{n, m})^p$, $0 \leq g_{n, m} \leq 1 < p$, and the sequence $g_{n, m}$ shall be integrable termwise in R .

COROLLARY 2.* *The double series of which the general term is $u_{n, m}(x, y)$ is absolutely convergent and absolutely integrable termwise in R if there exists a never negative function $v(x, y)$ which is integrable in R and a convergent double series of positive terms of the type $a_{n, m}$ such that $|u_{n, m}(x, y)| \leq a_{n, m}v(x, y)$.*

For, if $s_{n, m}$ refer to the sum of the absolute values of the nm terms of the u series, and $\sigma_{n, m}$ to the sum of the corresponding terms of the a series, $0 \leq s_{n, m} \leq \sigma_{n, m}v(x, y) \leq \lim_{n, m} \sigma_{n, m}v(x, y)$, and (d) of Corollary 1 applies.

* Cf. W. H. Young, *loc. cit.*, p. 324. The same statement holds good if R is a finite field.

8. Differentiation and Integration. The next theorem may be derived from what precedes in much the same manner as may be proved the theorem* of which it is a generalization.

THEOREM 13. Let $g_n(x, y) \leq f_n(x, y) \leq G_n(x, y)$, and $f_n - g_n$, $G_n - f_n$ be monotonic increasing functions of n at each point (x, y) , and let the derivative of $J_i(g_n) = J_i(dg_n/dn)$, and the derivative of $J_i(G_n) = J_i(dG_n/dn)$. Then the derivative with respect to n of $J_i(f_n) = J_i(df_n/dn)$ wherever $\dagger df_n/dn$ exists. The same is true if the right (left) hand derivative be used throughout instead of the derivative.

The following lemma may be proved by the use of the theorem of Young analogous to Theorem 9, and the theorem of Fubini analogous to the corollary of Theorem 5.

LEMMA. Let the interval $(a \leq n \leq b)$ be finite. Sufficient conditions that the following exist and be equal,

$$\int_a^b J_1[g(x, y; n)] dn, \quad J_1 \left[\int_a^b g(x, y; n) dn \right],$$

are that (1°) the integral of g with respect to x and y over $R_{\alpha, \beta}$, for all values of $\alpha, \beta > \alpha_0$, be absolutely less than an integrable function of n , (2°) $J_1(g)$ exist for each n , and g be absolutely integrable with respect to x, y , and n in the parallelopiped, $0 \leq x \leq \alpha$, $0 \leq y \leq \beta$, $a \leq n \leq b$.

THEOREM 14. Sufficient conditions that the equation,

$$\int_a^n J_1 \left(\frac{df}{dn} \right) dn = J_1(f_n) - J_1(f_a),$$

be valid in $(a \leq n \leq b)$ are that $J_1(f)$ exist at one point of the interval (a, b) , and that df/dn exist and satisfy the conditions imposed on g in the lemma.

By the lemma

$$\int_a^n J_1 \left(\frac{df}{dn} \right) dn = J_1 \int_a^n \frac{df}{dn} dn.$$

By a known theorem due to Vallée-Poussin, \ddagger this equals $J_1(f_n) - J_1(f_a)$.

COROLLARY. \S Under the same conditions the derivative, with respect to n , of $J_1(f_n)$ equals J_1 of the derivative of f_n , (a) in general in (a, b) including all

* See preceding note, same reference, p. 321.

\dagger More generally, for each value of n for which, except perhaps for a null set in the xy -plane, df/dn exists.

\ddagger *Loc. cit.*, Vol. I, p. 263.

\S See reference of previous note, pp. 267, 268.

points where the derivative of $J_1(f_n)$ exists, and (b) at all points of (a, b) provided $J_1(df/dn)$ is continuous in (a, b) .

THE FOURIER INTEGRALS, §§ 9–11.

9. Introduction. The iterated integral

$$J_4 = \frac{1}{\pi} \int_0^\infty du \int_{-\infty}^\infty f(t) \cos u(t-x) dt$$

is sometimes called the Fourier double integral. We shall reserve this name for

$$J_1 = D = \lim_{\alpha, \beta = \infty} \frac{1}{\pi} \int_0^\alpha du \int_0^\beta f(t) \cos u(t-x) dt + \lim_{\alpha, \beta = \infty} \frac{1}{\pi} \int_0^\alpha du \int_{-\beta}^0 f(t) \cos u(t-x) dt.$$

Here J_1 and J_4 have the same relative significance as in the preceding sections. It is evident that J_5 does not exist. A good deal has been written* recently on the problem of finding the most general conditions that one can impose on $f(t)$ and make J_4 equal to $f(x)$. In the following section, it is shown that J_1 equals $f(x)$ under circumstances more general than those derived for J_4 . Indeed, if $f(x) = 1$, J_4 does not exist, and if a value be given it by the usual summation method, that value is zero, but $J_1 = 1$. In Section 11 the corresponding integral which arises from three-dimensional problems is considered, and some of the questions connected with it. For simplicity x , y , and z are restricted to non-negative values. This enables one to dispense with the second term of D . No generality is lost thereby.

10. The Fourier Double Integral. LEMMA. (a) For all $\alpha, \beta \geq 0$ uniformly,

$$\left| \int_0^\alpha \frac{\sin \beta t}{t} dt \right| \leq \frac{\pi}{2}.$$

(b) Let $0 \leq f(t+x) \leq M$, and let f be a monotonic decreasing function of t ($t \geq 0$) for each x . Then, uniformly with respect to $\alpha \geq \alpha_0 > 0$, and boundedly† with respect to x in the first case, and uniformly in the second:

$$\lim_{\beta = \infty} \int_0^\alpha f(t+x) \frac{\sin \beta t}{t} dt = \frac{\pi}{2} f(+0+x), \quad \lim_{\beta = \infty} \int_{\alpha_0}^\alpha f(t+x) \frac{\sin \beta t}{t} dt = 0.$$

(c) If also uniformly with respect to x $\lim_{t=0} f(t+x) = f(+0+x)$, the convergence in the first case under (b) is uniform.

* A. Pringsheim, *Mathematische Annalen*, Vol. LXVIII (1910), pp. 367–408. W. H. Young, *Transactions of the Cambridge Philosophical Society*, Vol. XXI (1910), pp. 428–451; *Proceedings of the Royal Society of Edinburgh*, Vol. XXXI (1911), pp. 559–586.

† $F_n(x, y, \dots)$ approaches its limit “boundedly” with respect to x, y, \dots if there exists a fixed number greater than $|F_n|$ uniformly for all values of x, y, \dots .

The lemma is known except perhaps for the boundedness and uniformity of the convergence. By the more familiar part, however, we have

$$\begin{aligned} \frac{\pi}{2} f(+0+x) - \lim_{\beta=\infty} \int_0^a f(t+x) \frac{\sin \beta t}{t} dt \\ = \lim_{\beta=\infty} \int_0^a [f(+0+x) - f(t+x)] \frac{\sin \beta t}{t} dt = \lim_{\beta=\infty} S. \end{aligned}$$

By (a) and the second theorem of the mean,

$$|S| \leq M \max_a \left| \int_0^a \frac{\sin \beta t}{t} dt \right| \leq \frac{\pi M}{2}.$$

Thus the convergence is bounded. In the cases where it is uniform the demonstration is the same except that M is to be chosen arbitrarily.

THEOREM 15. *Let $0 \leq f(x) \leq M$, and let $f(x)$ be a monotonic decreasing function of $x(x \geq 0)$. Then $D = 1/2[f(x-0) + f(x+0)]$, and the proper integral of which D is the limit approaches D boundedly with respect to x if $x \geq a > 0$.*

CASE 1: ($x=0$). By the lemma

$$\pi D = \lim_{a, \beta=\infty} \int_0^a du \int_0^\beta f(t) \cos tu dt = \lim_{a, \beta=\infty} \int_0^\beta f(t) \frac{\sin at}{t} dt = \frac{\pi}{2} f(+0).$$

CASE 2: ($0 < a \leq x$). In the formal expression for D substitute $t-x=s$. Then

$$\begin{aligned} \pi D &= \lim_{a, \beta=\infty} \int_0^a du \int_{-x}^{\beta-x} f(s+x) \cos su ds = \lim_{a, \beta=\infty} \int_{-x}^{\beta-x} f(s+x) \frac{\sin as}{s} ds \\ &= \lim_{a, \beta=\infty} \int_{-x}^0 f(s+x) \frac{\sin as}{s} ds + \lim_{a, \beta=\infty} \int_0^{\beta-x} f(s+x) \frac{\sin as}{s} ds. \end{aligned}$$

Substituting $t=-s$ in the first integral, and separating the last integral into two parts by the point x ,

$$\pi D = \lim_{a, \beta=\infty} \int_0^x [f(-t+x) + f(t+x)] \frac{\sin at}{t} dt + \lim_{a, \beta=\infty} \int_x^{\beta-x} f(s+x) \frac{\sin as}{s} ds,$$

which equals $\frac{1}{2}[f(x-0) + f(x+0)]$ by (b) of the lemma, and the convergence is bounded.

COROLLARY 1. *If also $f(x)$ is continuous in $(0 < a \leq x \leq b)$, the convergence is uniform there.*

Apply (c) of the lemma.

COROLLARY 2. *The theorem and corollary are true if $f(x) = f_1(x) - f_2(x)$, where $f_1(x)$ and $f_2(x)$ have the properties there ascribed to $f(x)$.*

Examples: e^{-x} , $1 - e^{-x}$, $f(x)$ equal to a polynomial in $(0, 1)$ and to $1/x$ elsewhere.

THEOREM 16. Let (1°) $f(x)=g(x)p(x)$, where $0 \leq f, g, p \leq M$, $f(x)$ has limited variation in every finite* interval, and $g(x)$ is monotonic decreasing ($x \geq x_0$), and $p(x)$ is periodic ($x \geq x_0$). Let (2°), except perhaps a null set, for $x \geq x_0$,

$$p(x) = \sum_{i=0}^{\infty} (a_i \sin kix + b_i \cos kix),$$

where the a_i 's and b_i 's are the usual Fourier coefficients, and (3°) $\Sigma(|a_i| + |b_i|) = H$. Then for each $x \geq 0$, $D = 1/2[f(x-0) + f(x+0)]$.

Let $f(t) = \phi(t) + \psi(t)$, where $\phi = f$ and $\psi = 0$ at points where $t \leq 2x$, and $\phi = 0$ and $\psi = f$ where $t > 2x$, x being the point at which the development is to be examined. Referring to the preceding corollary we see that it holds when ϕ replaces f . Referring to the proof of the previous theorem we see that, by formal processes,

$$\begin{aligned} \pi D - \pi/2[f(x-0) + f(x+0)] &= \lim_{\alpha, \beta \rightarrow \infty} \int_0^x [\psi(-t+x) + \psi(t+x)] \frac{\sin \alpha t}{t} dt \\ &\quad + \lim_{\alpha, \beta \rightarrow \infty} \int_x^{\beta-x} \psi(s+x) \frac{\sin \alpha s}{s} ds = \text{I} + \text{II}. \end{aligned}$$

By definition of ψ , $\text{I} = 0$. To prove that $\text{II} = 0$ we have essentially to establish for the f of our present theorem only the second part of (b) of the lemma to Theorem 15, with the omission of the references to boundedness and uniformity with respect to x . An apparent exception occurs when $x = 0$, but it may be avoided by separating the interval (α, β) into two parts by means of a fixed point. We proceed, therefore, to show that, $\alpha_0 > 0$, $\epsilon > 0$, x being prescribed, there exists a β_ϵ so that the following is true uniformly in α and β :

$$\left| \int_{\alpha_0}^{\alpha} g(t+x) p(t+x) \frac{\sin \beta t}{t} dt \right| < \epsilon, \quad (\beta > \beta_\epsilon).$$

The function $g(t+x)/t$ is monotonic decreasing if $t > x_0$. Let α_1 be so large that $\alpha_1 > x_0$ and also that $g(t+x)/\alpha_1 < \epsilon$. Since $\dagger gp/t$ is absolutely L -integrable in (α_0, α_1) ,

$$\lim_{\beta \rightarrow \infty} \int_{\alpha_0}^{\alpha_1} \frac{gp}{t} \sin \beta t dt = 0.$$

Now

$$\begin{aligned} \left| \int_{\alpha_1}^{\alpha} \frac{g(t+x)}{t} p(t+x) \sin \beta t dt \right| &< \epsilon \max_{\lambda} \left| \int_{\alpha_1}^{\lambda} p(t+x) \sin \beta t dt \right| \\ &= \epsilon \max_{\lambda} \left| \int_{\alpha_1}^{\lambda} \sin \beta t \left[\sum_0^{\infty} a_i \sin ki(t+x) + b_i \cos ki(t+x) \right] dt \right|. \end{aligned}$$

* These conditions could be generalized a little by requiring that certain of them should hold necessarily only in the neighborhood of x .

† Lebesgue, *Annales de la Faculté de Toulouse*, Ser. 3, Vol. I (1909), p. 52.

It may be seen in several ways that this series is integrable termwise; for example, by 3° its remainder is uniformly limited. Thus the above expression equals

$$\varepsilon \max_{\lambda} \left| \sum_0^{\infty} \left[a_i \int_{a_1}^{\lambda} \sin ki(t+x) \sin \beta t dt + b_i \int_{a_1}^{\lambda} \cos ki(t+x) \sin \beta t dt \right] \right| \leq 8\varepsilon \Sigma(|a_i| + |b_i|) = 8\varepsilon H.$$

COROLLARY. *The theorem is true if $f(x) = f_1(x) - f_2(x)$, where f_1 and f_2 separately enjoy the properties there ascribed to f .*

Example: $f = x$ in $(0, \pi)$, $= 2\pi - x$ in $(\pi, 2\pi)$, etc.

THEOREM 17. *If $f(x)$ has limited variation in a neighborhood $(x-c, x+c)$, $c > 0$, of x , and if $f(x)/x$ is absolutely L -integrable in the infinite interval $(x \geq c)$, then, for each $x \geq 0$, $D = 1/2[f(x-0) + f(x+0)]$.*

As before

$$D = \lim_{a, \beta = \infty} \frac{1}{\pi} \int_{-x}^{\beta-x} f(t+x) \frac{\sin \alpha t}{t} dt = \lim_{a, \beta = \infty} \frac{1}{\pi} \left[\int_{-x}^{-\frac{c}{2}} \frac{f \sin \alpha t}{t} dt + \int_{-\frac{c}{2}}^{\frac{c}{2}} \frac{f \sin \alpha t}{t} dt + \int_{\frac{c}{2}}^{\beta-x} \frac{f \sin \alpha t}{t} dt \right] = \text{I} + \text{II} + \text{III}.$$

$$\text{II} = \lim_{a, \beta = \infty} \frac{1}{\pi} \int_0^{\frac{c}{2}} \frac{1}{2} [f(-t+x) + f(t+x)] \frac{\sin \alpha t}{t} dt = \frac{1}{2} [f(x-0) + f(x+0)],$$

since the part of the integrand in brackets has limited variation in $(0, c/2)$. Since $f(t+x)/t$ is absolutely L -integrable in $(-x, -c/2)$, by Lebesgue's* theorem I=0; and, similarly, a part of III,

$$\lim_{a, \beta = \infty} \frac{1}{\pi} \int_{\frac{c}{2}}^{\beta-x} f(t+x) \frac{\sin \alpha t}{t} dt = 0,$$

where β_{ϵ} is so chosen that

$$\int_{\beta_{\epsilon}}^{\infty} \left| \frac{f(t+x)}{t} \right| dt < \epsilon.$$

Thus $|\text{III}| < \epsilon$.

Example: $f(x) = p(x)e^{-x}$, where p is any limited function which has limited variation in the neighborhood of x and is L -integrable in every finite interval.

It could now be shown that, if in this form of the Fourier integral the integrand were multiplied by e^{-uy} (or by $e^{-c^2 u^2 t}$), the usual thermostatic (or thermodynamic) equations would be satisfied, in general, and that the functions

defined thereby would enjoy the other properties desired for the solutions of the corresponding problems in heat and electricity. But this would not be useful, for, although some texts imply that the Fourier "double" (in my sense, iterated) integral is needed in these physical problems, it never is, and the appropriate propositions can be proved for the simple integrals that are needed in a much more general fashion than I can prove them even for the double integrals considered here.* The theorems of this section serve as lemmas to the next section, where their practical value becomes apparent. However, as notably in the case of Fourier's series,† so to a less degree in the case of Fourier's integral, the theory of the representation of an arbitrary function has enjoyed a development and interest that have nothing to do with its value as a physical tool.

11. The Fourier Quadruple Integral. If one seeks the value of the potential function V in an infinite rectangular parallelopiped ($x, y, z \geq 0$), given that the surface $z=0$ is kept at a potential $V_0=f(x, y)$, and that the surface $x=0$ is insulated, one is led to LaPlace's equation, $\nabla^2(V)=0$, and to the formula: ‡

$$V_z = \lim_{\alpha, \beta, \gamma, \delta = \infty} \int_0^\alpha dt \int_0^\beta du \int_0^\gamma ds \int_0^\delta e^{-z\sqrt{s^2+u^2}} f(t, u) \cos s(t-x) \cos w(u-y) dw.$$

THEOREM 18. *Let $f(x, y)=h(x)k(y)$, where h and k satisfy the conditions imposed on f in one of the theorems 15, 16, or 17, or are the differences of two such functions. Then $V_0=1/4[f(x-0, y-0)+f(x-0, y+0)+f(x+0, y-0)+f(x+0, y+0)]$.*

For $V_0 = \lim_{\alpha, \beta, \gamma, \delta} \left[\int_0^\alpha dt \int_0^\gamma h(t) \cos s(t-x) ds \right] \left[\int_0^\beta du \int_0^\delta k(y) \cos w(u-y) dw \right]$, which equals $1/4[h(x-0)+h(x+0)][k(y-0)+k(y+0)]$.

COROLLARY. *The convergence is uniform with respect to x (or y) if h (or k) also satisfies the condition of continuity imposed in the corollary to Theorem 15.*

THEOREM 19. *If $f(x, y)$ is absolutely L -integrable in the infinite rectangle ($x, y \geq 0$), $V_z(z>0)$ satisfies LaPlace's equation.*

It is sufficient to show that the signs of partial differentiation may be passed over the limit sign and over the signs of integration in the expression for V_z . For this purpose we refer to corollary (b) of Theorem 14. Using

* Cf. W. H. Young, *Proceedings of the Royal Society of Edinburgh*, Vol. XXXI (1911), p. 587; *Encyklopädie Mathematischen Wissenschaften*, II, A 12, p. 1089.

† Cf. VanVleck, "The Influence of Fourier's Series upon the Development of Mathematics," *Science*, N. S., Vol. XXXIX (1914), Number 995, pp. 113-124.

‡ In the usual formula the iterated integral is used.

the notation there, but replacing that f by the present integrand, which we call ϕ , and thinking of n as representing in turn x , y , and z , we see that we need to prove: (1) that $J_1(d\phi/dn)$ is continuous near n , n being the point at which the derivative of J_1 is being examined; (2) that $J_1(\phi)$ exists at n ; (3) that $d\phi/dn$ exists, and is absolutely L -integrable over finite three-dimensional fields; and (4) that there exists a function, $\Phi(n)$, whose integral with respect to n exists over a finite interval enclosing the particular point n in question, such that, for all values of $\alpha, \beta, \gamma, \delta \geq 0$ uniformly,

$$\left| \int_{R_{\alpha, \beta, \gamma, \delta}} \frac{d\phi}{dn} dt du ds dw \right| < \Phi(n).$$

These results are needed in order to show that the first derivative of J_1 may be replaced by the first derivative of its integrand. The process must be repeated for the second derivative. We shall then have conditions (5), (6), (7), and (8) exactly like (1), (2), (3), and (4), respectively, except that $d\phi/dn$ replaces ϕ , and in (4) $\Phi(n)$ is replaced by some other function having the same properties. It is obvious that (3) and (7) are true, and that (6) is included in (1). We first prove (4). For all values of x, y, z uniformly,

$$|\phi| \leq |f(t, u)| e^{-z\sqrt{s^2+w^2}} \leq |f| e^{-z(\frac{s+w}{2})} = |f| e^{-\frac{zs}{2}} e^{-\frac{zw}{2}},$$

$$\left| \int_{R_{\alpha, \beta, \gamma, \delta}} \phi \right| \leq M \left(\int_0^\infty e^{-\frac{zs}{2}} ds \right)^2 = \frac{4M}{z^2},$$

where

$$M = \lim_{\alpha, \beta} \int_0^\alpha dt \int_0^\beta |f(t, u)| du.$$

Thus (2) $J_1(\phi)$ exists absolutely. Now, first let $n=z$.

$$\left. \begin{aligned} \frac{\partial \phi}{\partial z} &= -\sqrt{s^2+w^2}\phi, \quad \frac{\partial^2 \phi}{\partial z^2} = (s^2+w^2)\phi, \quad \left| \frac{\partial \phi}{\partial z} \right| \leq |f|(s+w)e^{-\frac{zs}{2}}e^{-\frac{zw}{2}}, \\ \left| \int_{R_{\alpha, \beta, \gamma, \delta}} \frac{\partial \phi}{\partial z} \right| &\leq 2M \left(\int_0^\infty s e^{-\frac{zs}{2}} ds \int_0^\infty e^{-\frac{zw}{2}} dw \right) = \frac{16M}{z^3}. \end{aligned} \right\} \quad (9)$$

Thus (4) is satisfied ($z > 0$). We can similarly prove (8), for

$$\left| \int_{R_{\alpha, \beta, \gamma, \delta}} \frac{\partial^2 \phi}{\partial z^2} \right| \leq 2M \int_0^\infty s^2 e^{-\frac{zs}{2}} ds \int_0^\infty e^{-\frac{zw}{2}} dw = \frac{64M}{z^4}.$$

Now, by referring to Corollary 1 (d°) of Theorem 12, it will be seen that in (9) we have established (1) and (5); *e. g.*, if $z \geq c > 0$,

$$|f|(s+w)e^{-\frac{z}{2}(s+w)} \leq |f|(s+w)e^{-\frac{c}{2}(s+w)},$$

which is an integrable function in R . Secondly, let $n=x \geq 0$.

$$\frac{\partial \phi}{\partial x} = s e^{-z\sqrt{s^2+w^2}} f(t, u) \sin s(t-x) \cos w(u-y),$$

$$\frac{\partial^2 \phi}{\partial x^2} = -s^2 e^{-z\sqrt{s^2+w^2}} \cos s(t-x) \cos w(u-y).$$

$$\left| \int_{R_{\alpha, \beta, \gamma, \delta}} \frac{\partial \phi}{\partial x} \right| < M \int_0^\infty s e^{-\frac{zs}{2}} ds \int_0^\infty e^{-\frac{zw}{2}} dw = \frac{8M}{2^3}.$$

$$\left| \int_{R_{\alpha, \beta, \gamma, \delta}} \frac{\partial^2 \phi}{\partial x^2} \right| < M \int_0^\infty s^2 e^{-\frac{zs}{2}} ds \int_0^\infty e^{-\frac{zw}{2}} dw = \frac{32M}{2^4}.$$

Hence the conditions in question are satisfied. The case, $n=y \geq 0$, is identical with this one.

WESLEYAN UNIVERSITY, MIDDLETOWN, CONN., *October, 1916.*